

CMS Search Plans and Sensitivity to New Physics with Dijets

The CMS Collaboration

Abstract

1 CMS will use dijets to search for physics beyond the standard model during early LHC
2 running. The inclusive jet cross section as a function of jet transverse momentum with
3 10 pb^{-1} of integrated luminosity is sensitive to contact interactions beyond the reach of
4 the Tevatron. The dijet mass distribution will be used to search for dijet resonances such
5 as an excited quark. Additional sensitivity to the existence of contact interactions or
6 dijet resonances can be obtained by comparing dijet rates in two distinct pseudorapidity
7 regions.

The Large Hadron Collider at CERN will produce many events with two energetic jets resulting from proton-proton collisions at $\sqrt{s} = 14$ TeV. These dijet events result from parton scattering, produced by the strong interaction of quarks (q) and gluons (g) inside the protons. This paper discusses plans to use dijets in the search for two signals of new physics: contact interactions and resonances decaying into dijets. This generic search is applied to two models of quark compositeness, that are used as benchmarks of sensitivity to new physics. The first model is a contact interaction [1] among left-handed quarks at an energy scale Λ^+ in the process $qq \rightarrow qq$, modeled with the effective Lagrangian $L_{qq} = (\pm 2\pi\alpha_s/\Lambda^2)(\bar{q}_L\gamma^\mu q_L)(\bar{q}_L\gamma_\mu q_L)$ with $+$ chosen for the sign. The second model is a dijet resonance signal from the decay of an excited quark (q^*) [2] in the process $qg \rightarrow q^* \rightarrow qq$. All processes presented here have been simulated using PYTHIA version 6.4 [3].

A detailed description of the Compact Muon Solenoid (CMS) experiment can be found elsewhere [4, 5]. The CMS coordinate system has the origin at the center of the detector, z -axis points along the beam direction toward the Jura mountains, transverse direction perpendicular to the beam, azimuthal angle ϕ , polar angle θ , and pseudorapidity $\eta = -\ln(\tan[\theta/2])$. The central feature of the CMS apparatus is a superconducting solenoid, of 6m internal diameter. Within the field volume are the silicon pixel and strip tracker, and the barrel and endcap calorimeters. The crystal electromagnetic calorimeter (ECAL) and the brass-scintillator hadronic calorimeter (HCAL) are in the barrel and endcap ($|\eta| < 3$). Outside the field volume in the forward region is an iron-quartz fiber hadronic calorimeter

($3 < |\eta| < 5$). The HCAL and ECAL cells are grouped into towers, projecting radially outward from the origin, for triggering purposes and to facilitate the jet reconstruction. In the region $|\eta| < 1.74$ these projective calorimeter towers have segmentation $\Delta\eta = \Delta\phi = 0.087$, and the η and ϕ width progressively increases at higher values of η . The energy in the HCAL and ECAL within each projective tower is summed to find the calorimeter tower energy. Towers with $|\eta| < 1.3$ contain only cells from the barrel calorimeters, towers in the transition region $1.3 < |\eta| < 1.5$ contain a mixture of barrel and endcap cells, and towers in the region $1.5 < |\eta| < 3.0$ contain only cells from the endcap calorimeters.

Jets are reconstructed using both the iterative and midpoint cone algorithms [5], with indistinguishable results for this analysis. Below we will discuss three types of jets: reconstructed, corrected and generated. The reconstructed jet energy, E , is defined as the scalar sum of the calorimeter tower energies inside a cone of radius $\sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.5$, centered on the jet axis. The jet momentum, \vec{p} , is the corresponding vector sum: $\vec{p} = \sum E_i \hat{u}_i$ with \hat{u}_i being the unit vector pointing from the origin to the energy deposition E_i inside the same cone. Jet transverse momentum, p_T , is the component of \vec{p} in the transverse direction. The E and \vec{p} of a reconstructed jet are then corrected for the non-linear response of the calorimeter to a generated jet. Generated jets come from applying the same jet algorithm to the Lorentz vectors of stable generated particles before detector simulation. The p_T of a corrected jet is equal to the p_T of the corresponding generated jet on average. The jet corrections estimated from a GEANT [6] simulation of the CMS detector increase the jet p_T on average by roughly 50% (10%) for 70 GeV (3 TeV) jets

in the region $|\eta| < 1.3$. Further details on jet reconstruction and jet energy corrections can be found elsewhere [5, 7]. The jet measurements presented here are within the region $|\eta| < 1.3$, where there is expected to be the highest sensitivity to new physics, and where the reconstructed jet response variations as a function of η are both moderate and smooth.

The dijet system is defined to be the two jets with the highest p_T in an event (leading jets), and the dijet mass is given by $m = \sqrt{(E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2}$. The estimated dijet mass resolution varies from 9% at a dijet mass of 0.7 TeV to 4.5% at 5 TeV.

CMS will record events that pass a first level trigger and a high level trigger. For an instantaneous luminosity of $10^{32} \text{ cm}^{-2}\text{s}^{-1}$, consider three event samples collected by requiring at least one jet in the high level trigger with corrected transverse energy above 60, 120 and 250 GeV, prescaled by factors of 2000, 40 and 1, respectively. For an integrated luminosity of 100 pb^{-1} , the three event samples will effectively correspond to 0.05, 2.5, and 100 pb^{-1} . The first event sample will be used to measure the trigger efficiency of the second sample. The last two event samples will be used to study dijets of mass above 330 and 670 GeV, respectively, where the trigger efficiencies are expected to be higher than 99% [9].

Backgrounds from cosmic-rays, beam halo, and detector noise are expected to occasionally produce events with large or unbalanced energy depositions. They will be removed by requiring $\cancel{E}_T / \sum E_T < 0.3$ and $\sum E_T$ less than 14 TeV, where \cancel{E}_T ($\sum E_T$) is the magnitude of the vector (scalar) sum of the transverse energies measured by all calorimeter towers in the event. This cut is estimated to be more than 99% efficient for both QCD

jet events and the signals of new physics considered. In the high p_T region relevant for this search, jet reconstruction is fully efficient.

CMS plans to search for contact interactions using the jet p_T distribution. Figure 1 shows the inclusive jet differential cross section as a function of p_T , for jets with $|\eta| < 1$. Considering first the QCD processes, the reconstructed and corrected quantities are compared with the QCD prediction for generated jets. After corrections, the reconstructed and generated distributions agree. The ratio of the corrected jet cross section to the generated jet cross section varies between 1.2 at $p_T = 100$ GeV and 1.05 at $p_T = 500$ GeV, and remains roughly constant for higher p_T . The deviation of this ratio from 1 is attributed to the smearing effect of the jet p_T resolution on the steeply falling spectrum. The measured spectrum could be further corrected for resolution smearing, and this ratio from Monte Carlo is an estimate of the size of that correction. The measurement uncertainties are predominantly systematic. The inset in figure 1 shows the effect on the jet rate of a 10% uncertainty in jet energy correction that could be expected in early running when only 10 pb^{-1} of integrated luminosity have been accumulated. This experimental uncertainty is roughly ten times larger than the uncertainties from parton distributions, as estimated using CTEQ6.1 fits [10]. Figure 1 shows that the effect of new physics from a contact interaction with scale $\Lambda^+ = 3 \text{ TeV}$ is convincingly above what could be expected for measurement uncertainties with only 10 pb^{-1} . For comparison, a Tevatron search has excluded contact interactions with scales Λ^+ below 2.7 TeV [11].

CMS plans to search for narrow dijet resonances using the dijet mass distribution.

92 Figure 2 shows the differential cross section versus dijet mass, where both leading jets
 93 have $|\eta| < 1$, and the mass bins have a width roughly equal to the dijet mass resolution.
 94 Considering first the QCD processes, the cross section for corrected jets agrees with the
 95 QCD prediction from generated jets. To determine the background shape either the
 96 Monte Carlo prediction or a parameterized fit to the data can be used. The inset to
 97 figure 2 shows a simulation of narrow dijet resonances with a q^* production cross section.
 98 This is compared to the QCD statistical uncertainties including trigger prescaling. This
 99 shows that with an integrated luminosity of 100 pb^{-1} a q^* dijet resonance with a mass
 100 of 2 TeV would produce a convincing signal above the statistical uncertainties from the
 101 QCD background. For comparison, a Tevatron search has excluded q^* dijet resonances
 102 with mass, M , below 0.87 TeV [12]. The heaviest dijet resonances that CMS can discover
 103 (at five standard deviations) with 100 pb^{-1} of integrated luminosity, using this search
 104 technique and including the expected systematic uncertainties [13, 14], are: 2.5 TeV for
 105 q^* , 2.2 TeV for axigluons [15] or colorons [16], 2.0 TeV for E_6 diquarks [17], and 1.5 TeV
 106 for color octet technirhos [18]. Studies of the jet η cut have concluded that the optimal
 107 sensitivity to new physics is achieved with $|\eta| < 1.3$ for a 2 TeV spin 1 dijet resonance
 108 decaying to $q\bar{q}$ [8].

109 CMS plans to search for both contact interactions and dijet resonances using the dijet
 110 ratio, $r = N(|\eta| < 0.7)/N(0.7 < |\eta| < 1.3)$, where N is the number of events with both jets
 111 in the specified $|\eta|$ region. The dijet ratio is sensitive to the dijet angular distribution. For
 112 the QCD processes, the dijet ratio is the same for corrected jets and generated jets, and

is constant at $r = 0.5$ for dijet masses up to 6 TeV [8]. Figure 3 shows the dijet ratio from contact interactions and dijet resonances, compared to the expected statistical uncertainty on the QCD processes, for 100 pb⁻¹ of integrated luminosity, including trigger prescaling. The signal from a contact interaction with scale $\Lambda^+ = 5$ TeV rises well above the QCD statistical errors at high dijet mass. Systematic uncertainties in the dijet ratio are expected to be small, since they predominantly cancel in the ratio as previously reported [13, 19]. Using the dijet ratio, CMS can discover a contact interaction at scale $\Lambda^+ = 4, 7$ and 10 TeV with integrated luminosities of 10, 100, and 1000 pb⁻¹, respectively [8]. The signal from a 2 TeV spin 1/2 q^* produces a convincing peak in the dijet ratio, because it has a significant rate and a relatively isotropic angular distribution compared to the QCD t -channel processes. Fixing the cross section of the 2 TeV dijet resonance for $|\eta| < 1.3$ at 13.6 pb (from the q^* model), the dijet ratio in the presence of QCD background increases by approximately 6% when considering a spin 2 resonance decaying to both $q\bar{q}$ and gg (such as a Randall-Sundrum graviton [20]), and the dijet ratio decreases by approximately 4% when considering a spin 1 resonance decaying to $q\bar{q}$ (such as a Z' , axigluon, or coloron) [8]. Hence, the sensitivity to a 2 TeV dijet resonance depends only weakly on the spin of the resonance. Nevertheless, with sufficient luminosity, this simple measure of the dijet angular distribution, or a more complete evaluation of the angular distribution, can be used to see these small variations and infer the spin of a dijet resonance.

In conclusion, CMS plans to use measurements of rate as a function of jet p_T and dijet

mass, as well as a ratio of dijet rates in different η regions, to search for new physics in the data sample collected during early LHC running. With integrated luminosity samples in the range 10–100 pb⁻¹, CMS will be sensitive to contact interactions and dijet resonances beyond those currently excluded by the Tevatron.

We thank the staffs at CERN and other CMS Institutes, and acknowledge support from: Austrian Federal Ministry of Science and Research; FNRS and FWO (Belgium); CNPq and FAPERJ (Brazil); Bulgarian Ministry of Education and Science; CERN; CAS and NSFC (China); Croatian Ministry of Science and Technology; University of Cyprus; Estonian Academy of Sciences and NICPB; Academy of Finland, Finish Ministry of Education and Helsinki Institute of Physics; CEA and CNRS/IN2P3 (France); BMBF and DESY (Germany); General Secretariat for Research and Technology (Greece); NKTH (Hungary); DAE and DST (India); IPM (Iran); UCD (Ireland); INFN (Italy); KICOS (Korea); CINVESTAV, CONACYT, and UASLP-FAI (Mexico); PAEC (Pakistan); State Commission for Scientific Research (Poland); FCT (Portugal); JINR (Armenia, Belarus, Georgia, Ukraine, Uzbekistan), Ministry of Science and Technologies of the Russian Federation, Russian Ministry of Atomic Energy; Ministry of Science and Environmental Protection of the Republic of Serbia; Oficina de Ciencia y Tecnologia (Spain); ETHZ, PSI, University of Zurich (Switzerland); National Science Council (Taipei); TAEA (Turkey); STFC (United Kingdom); DOE and NSF (USA).

References

- [1] E. Eichten, K. Lane and M. E. Peskin, Phys. Rev. Lett. **50**, 811 (1983).
- [2] U. Baur, I. Hinchliffe and D. Zeppenfeld, Int. J. Mod. Phys. **A2**, 1285 (1987).
- [3] T. Sjostrand, L. Lonnblad, S. Mrenna, hep-ph/0108264.
- [4] CMS Collaboration, “The CMS experiment at the CERN LHC” (accepted for publication in JINST).
- [5] CMS Collaboration, Physics TDR Volume I, CERN-LHCC-2006-001.
- [6] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Meth. **A506**, 250-303 (2003).
- [7] CMS Collaboration, CMS Physics Analysis Summary JME-07-003 (2008). at <http://cms-physics.web.cern.ch/cms-physics/public/JME-07-003-pas.pdf> .
- [8] CMS Collaboration, CMS Physics Analysis Summary SBM-07-001 (2007) at <http://cms-physics.web.cern.ch/cms-physics/public/SBM-07-001-pas-v3.pdf> .
- [9] S. Esen and R. Harris, CMS Note 2006/069.
- [10] J. Pumplin *et al.* (CTEQ Collaboration), JHEP **07**, 012 (2002).
- [11] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. Lett. **82**, 2457 (1999).
- [12] K. Hatakeyama *et al.* (CDF Collaboration) recent public results at http://www-cdf.fnal.gov/physics/exotic/r2a/20080214.mjj_resonance_1b/.

- 171 [13] CMS Collaboration, J. Phys. G: Nucl. Part. Phys. **34**, 995 (2006).
- 172 [14] K. Gumus, N. Akchurin, S. Esen and R. Harris, CMS Note 2006/070.
- 173 [15] J. Bagger, C. Schmidt and S. King, Phys. Rev. D**37**, 1188 (1988).
- 174 [16] E. Simmons, Phys. Rev. D**55**, 1678 (1997).
- 175 [17] J. Hewett and T. Rizzo, Phys. Rep. **183**, 193 (1989) and references therein.
- 176 [18] K. Lane and S. Mrenna, Phys. Rev. D**67**, 115011 (2003).
- 177 [19] S. Esen and R. Harris, CMS Note 2006/071.
- 178 [20] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).




Figure 1: The inclusive jet p_T differential cross section expected from QCD for $|\eta| < 1$, for generated jets (points), reconstructed jets (triangles), and corrected jets (open circles). The inset shows the number of generated jets expected in 50 GeV bins for an integrated luminosity of 10 pb^{-1} . The standard QCD processes (solid) are compared to a signal from contact interactions with scale $\Lambda^+ = 3 \text{ TeV}$ (dotted) and 5 TeV (dashed). The shaded band represents the effect of a 10% uncertainty on the jet energy scale.




Figure 2: The dijet mass differential cross section expected from QCD for $|\eta| < 1$ vs. dijet mass, m , for generated jets (points), reconstructed jets (triangles), and corrected jets (open boxes). The inset shows dijet resonances reconstructed using corrected jets coming from q^* signals [14] of mass 0.7, 2, and 5 TeV. The fractional difference (histogram) between the q^* signal and the QCD background is compared to the QCD statistical error (vertical bars) for an integrated luminosity of 100 pb^{-1} .




Figure 3: The dijet ratio for corrected jets expected from QCD (horizontal line), with statistical uncertainties (vertical bars) for an integrated luminosity of 100 pb^{-1} , is compared to QCD + contact interaction signals with a scale $\Lambda^+ = 5 \text{ TeV}$ (dashed) and 10 TeV (dotted), as well as to QCD + dijet resonance signals (histogram) with q^* masses of 0.7 and 2 TeV .